ONE MONTH TYPICAL DEVELOPMENT CHECKLIST (©Diane Bahr)

Place a check mark next to the characteristics you see in your one-month-old baby

Typical 1 Month Old Baby	YES	NO
FEEDING/MOUTH DEVELOPMENT		
Begins gaining control over rooting reflex (rooting leads to sucking)		
Locates nipple with mouth easily		
Sequences 2 or more sucks from the bottle or breast		
Mouths hands & fingers near front of mouth (generalized mouthing)		
Mouth, nose, & throat growing & changing (different from newborn)		
Has closed mouth & easy nose-breathing when mouth is inactive & empty		
Rests tongue in roof of the mouth to help maintain broad palate shape		
Is free of tethered tongue, lip, and/or cheek ties		
Can mirror/imitate some mouth movement		
Can match some pitch & duration of human voice		
BODY DEVELOPMENT		
When Held Upright		
Lifts & holds head up in line with body for a few seconds (birth to 1 month)		
Holds head up steadily in line with body for a few minutes (1 to 2 months)		
When on the Stomach		
Rests head to one side or the other & can bring hand to mouth on that side		
Moves arms & legs away from body & lowers hips toward a surface when resting		
Lifts & turns head with increasing skill (shifts weight to arms; shoulders likely sending		
information through spine for hips to lift from surface)		
Can lift head for a few moments without chin touching a surface (birth to 1 month)		
Can lift head for approximately 5 seconds (1 to 2 months)		
Moves legs in thrusting or crawling motion (birth to 1 month)		
When on Each Side		
Turns from side to back from each side (1 to 2 months)		
·		
When on the Back		
Has increased skill in head turning & chin lifting; may turn head from side-to-side		
May assume fencing posture with head turning (arm reaching outward on face side,		
elbow bent on skull side – asymmetric tonic neck reflex/response)		
May turn head to look at something & move hands in a rudimentary way (possible eye-		
hand connection)		
Has better peripheral than midline vision, so can track object from side to middle of body		
Has increased chest, shoulder, arm, & hand opening likely secondary to gravitational pull		
Closes hand around adult's finger or object (grasp response)		
Moves arms vigorously (1 to 2 months)		
Kick legs in the air reciprocally, as well as symmetrically		
Bends body, arms, & legs at the same time (tonic labyrinthine reflex, birth to 1 month)		

Straightens body; tries to straighten arms & legs (tonic labyrinthine reflex, birth to 1 month)

Primary References

- Bahr, D. (2018). Feed your baby and toddler right: Early eating and drinking skills encourage the best development. Arlington, TX: Future Horizons.
- Bahr, D. C., & Hillis, A. E. (2001). Neurological and anatomical bases. In D. C. Bahr *Oral motor assessment and treatment: Ages and stages* (pp. 1-41). USA: Pearson College Division.
- Bly, L. (1994). Motor skills acquisition in the first year: An illustrated guide to normal development. USA: Psychological Corp.
- Feldenkrais, M. (1972). Awareness through movement (Vol. 1977). New York: Harper and Row.
- Morris, S. E., & Klein, M. D. (2000). *Pre-Feeding skills: A comprehensive resource for mealtime development*. (2nd ed.). San Antonio, TX: Therapy Skill Builders.
- Neuro Restart (2019). Primitive reflexes. Retrieved from http://www.neurorestart.co.uk/primitive-reflexes/
- Vulpe, S. G. (1994). Vulpe assessment battery-revised: Developmental assessment, performance analysis, individualized programming for the atypical child. East Aurora, NY: Slosson Educational Publications.

Checklist User Information: This checklist is meant for educational purposes only. It is not meant to diagnose or treat any disorders or to provide health, medical, or professional consultation. Participants should always seek the advice of appropriate professionals as needed. Diane Bahr is not responsible for risk, loss, or liability as a direct or indirect consequence related to information found in this checklist. Your use of the checklist implies your agreement to and understanding of these terms.